

Problem 18) a) The slope of the function $y = ax^2$ at $x = x_0$, namely, $y' = 2ax_0$, is the tangent of the angle φ that the unit-vector \mathbf{S} makes with the x -axis. Therefore,

$$\tan \varphi = 2ax_0.$$

$$\cos \varphi = 1/\sqrt{1 + \tan^2 \varphi} = 1/\sqrt{1 + 4a^2x_0^2}.$$

$$\sin \varphi = \tan \varphi \cos \varphi = 2ax_0/\sqrt{1 + 4a^2x_0^2}.$$

$$\mathbf{S} = \cos \varphi \hat{\mathbf{x}} + \sin \varphi \hat{\mathbf{y}} = (\hat{\mathbf{x}} + 2ax_0\hat{\mathbf{y}})/\sqrt{1 + 4a^2x_0^2}.$$

$$\mathbf{R}_2 = [-x_0\hat{\mathbf{x}} + (y_1 - y_0)\hat{\mathbf{y}}]/\sqrt{x_0^2 + (y_1 - y_0)^2}.$$

b) The dot-product of \mathbf{S} and \mathbf{R}_1 gives the cosine of the angle between these two unit-vectors. Similarly, the cosine of the angle between \mathbf{S} and \mathbf{R}_2 is given by $\mathbf{S} \cdot \mathbf{R}_2$. Given that $\mathbf{R}_1 = -\hat{\mathbf{y}}$, and that \mathbf{R}_2 and \mathbf{S} are found in part (a), we now set $\mathbf{S} \cdot \mathbf{R}_1 = \mathbf{S} \cdot \mathbf{R}_2$. Consequently, the various system parameters are related as follows:

$$-2ax_0 = [-x_0 + 2ax_0(y_1 - y_0)]/\sqrt{x_0^2 + (y_1 - y_0)^2}.$$

c) The above equation may now be solved to determine y_1 in terms of the remaining parameters. Recalling that $y_0 = ax_0^2$, we will have

$$2a = [1 - 2a(y_1 - ax_0^2)]/\sqrt{x_0^2 + (y_1 - ax_0^2)^2}$$

$$\rightarrow 4a^2[x_0^2 + (y_1 - ax_0^2)^2] = 1 - 4a(y_1 - ax_0^2) + 4a^2(y_1 - ax_0^2)^2$$

$$\rightarrow 4a^2x_0^2 = 1 - 4ay_1 + 4a^2x_0^2 \quad \rightarrow \quad y_1 = 1/(4a).$$
