

Problem 2) a) $\mathcal{F}\{\cos(2\pi f_0 x)\} = \frac{1}{2} \mathcal{F}\{\exp(i2\pi f_0 x) + \exp(-i2\pi f_0 x)\} = \frac{1}{2}[\delta(s - f_0) + \delta(s + f_0)].$

b) $\mathcal{F}\{\sin(2\pi f_0 x)\} = \frac{1}{2i} \mathcal{F}\{\exp(i2\pi f_0 x) - \exp(-i2\pi f_0 x)\} = \frac{1}{2i}[\delta(s - f_0) - \delta(s + f_0)].$

c) $\mathcal{F}\{\cos^2(\pi f_0 x)\} = \frac{1}{2} \mathcal{F}\{1 + \cos(2\pi f_0 x)\} = \frac{1}{2}\delta(s) + \frac{1}{4}[\delta(s - f_0) + \delta(s + f_0)].$

d) Differentiation theorem:

$$f(x) = \int_{-\infty}^{\infty} F(s) \exp(i2\pi s x) ds \rightarrow \frac{df(x)}{dx} = \int_{-\infty}^{\infty} i2\pi s F(s) \exp(i2\pi s x) ds \rightarrow \mathcal{F}\left\{\frac{df(x)}{dx}\right\} = i2\pi s F(s).$$

Therefore,

$$\mathcal{F}\left\{\frac{d}{dx} \cos^2(\pi f_0 x)\right\} = i2\pi s \left\{\frac{1}{2}\delta(s) + \frac{1}{4}[\delta(s - f_0) + \delta(s + f_0)]\right\} = 0 + \frac{i2\pi}{4}[f_0\delta(s - f_0) - f_0\delta(s + f_0)].$$

Carrying out the differentiation, we find

$$\mathcal{F}\{-2\pi f_0 \sin(\pi f_0 x) \cos(\pi f_0 x)\} = -\pi f_0 \mathcal{F}\{\sin(2\pi f_0 x)\} = \frac{i\pi f_0}{2}[\delta(s - f_0) - \delta(s + f_0)].$$

Consequently,

$$\mathcal{F}\{\sin(2\pi f_0 x)\} = \frac{1}{2i}[\delta(s - f_0) - \delta(s + f_0)].$$

This is the same result as obtained in part (b).