

Quantum States of the Quantized Field


Amplitude and Phase

- Key characteristics of classical fields
- Need equivalents for quantum fields

Classical Field

$$E(z,t) = \mathcal{E}_0 \alpha e^{-i(\omega t - kz)} + \text{c.c.}$$

\uparrow
 $|\alpha| e^{i\varphi}$



Quantum Field

$$\hat{E}(z,t) = \mathcal{E}_0 \hat{a} e^{-i(\omega t - kz)} + \text{H.C.}$$

\uparrow
Non-Hermitian!

Separate in amplitude & phase?

Consider operators

$$\hat{a} = (\hat{N}+1)^{1/2} \hat{x}_p(i\varphi)$$

$$\hat{a}^\dagger = \hat{x}_p(-i\varphi) (\hat{N}+1)^{1/2}$$

“phase” “amplitude”



$$\hat{x}_p(i\varphi) = (\hat{N}+1)^{-1/2} \hat{a}$$

$$\hat{x}_p(-i\varphi) = \hat{a}^\dagger (\hat{N}+1)^{-1/2}$$

“Phase operators”

$$\hat{x}_p(i\varphi) \hat{x}_p(-i\varphi) = 1 \quad \hat{x}_p(i\varphi) = \hat{x}_p(-i\varphi)^\dagger$$

$$\hat{x}_p(-i\varphi) \hat{x}_p(i\varphi) = 1 \quad = [\hat{x}_p(-i\varphi)]^{-1}$$

- Analogous to classical phases
- Non-Hermitian, NOT observables

Quadrature operators?

$$\hat{c} \cos \varphi = \frac{1}{2} [\hat{x}_p(i\varphi) + \hat{x}_p(-i\varphi)]$$

$$= \frac{1}{2} [(\hat{N}+1)^{-1/2} \hat{a} + \hat{a}^\dagger (\hat{N}+1)^{-1/2}]$$

$$\hat{s} \sin \varphi = \frac{1}{2i} [\hat{x}_p(i\varphi) - \hat{x}_p(-i\varphi)]$$

$$= \frac{1}{2i} [(\hat{N}+1)^{-1/2} \hat{a} - \hat{a}^\dagger (\hat{N}+1)^{-1/2}]$$

- Hermitian -> observables
- but ultimately too cumbersome

Let's rewind and try again...

Quantum States of the Quantized Field

“Phase operators”

$$e^{\hat{x}p(i\varphi)} e^{\hat{x}p(-i\varphi)} = 1 \quad e^{\hat{x}p(i\varphi)} = e^{\hat{x}p(-i\varphi)^\dagger}$$

$$e^{\hat{x}p(-i\varphi)} e^{\hat{x}p(i\varphi)} = 1 \quad = [e^{\hat{x}p(-i\varphi)}]^{-1}$$

- Analogous to classical phases
- Non-Hermitian, NOT observables

Quadrature operators?


$$\begin{aligned} \hat{\cos}\varphi &= \frac{1}{2} [e^{\hat{x}p(i\varphi)} + e^{\hat{x}p(-i\varphi)}] \\ &= \frac{1}{2} [(\hat{N}+1)^{-1/2} \hat{a} + \hat{a}^\dagger (\hat{N}+1)^{-1/2}] \\ \hat{\sin}\varphi &= \frac{1}{2i} [e^{\hat{x}p(i\varphi)} - e^{\hat{x}p(-i\varphi)}] \\ &= \frac{1}{2i} [(\hat{N}+1)^{-1/2} \hat{a} - \hat{a}^\dagger (\hat{N}+1)^{-1/2}] \end{aligned}$$

- Hermitian → observables
- but ultimately too cumbersome

Let's rewind and try again...

Quadratures of the Classical Field – Take Two

$$E(z,t) = \sum_k \underbrace{\alpha_k(t)}_{\text{complex amplitude for mode } e^{ikz}} e^{ikz} + \text{c.c.}$$



Define

$$\begin{aligned} X(t) &= \text{Re}[\alpha_k(t)] = \frac{1}{2} [\alpha_k(t) + \alpha_k^*(t)] = Q(t) \\ Y(t) &= \text{Im}[\alpha_k(t)] = \frac{1}{2i} [\alpha_k(t) - \alpha_k^*(t)] = P(t) \end{aligned}$$

Quantization: $\alpha \rightarrow \hat{a}, \alpha^* \rightarrow \hat{a}^\dagger$

$$\left. \begin{aligned} \hat{X}(t) &= \frac{1}{2} [\hat{a}_k(t) + \hat{a}_k^\dagger(t)] = \hat{Q}(t) \\ \hat{Y}(t) &= \frac{1}{2i} [\hat{a}_k(t) - \hat{a}_k^\dagger(t)] = \hat{P}(t) \end{aligned} \right\} [\hat{X}(t), \hat{Y}(t)] = i/2$$

$$\begin{aligned} \hat{E}(z,t) &= \sum_k (\hat{X}(t) + i\hat{Y}(t)) e^{ikz} + \text{H.C.} \\ &= \sum_k [\hat{X}(t) \cos(kz) - \hat{Y}(t) \sin(kz)] \end{aligned}$$


– same info, easier to work with –

Quantum States of the Quantized Field

Quadratures of the Classical Field – Take Two

$$E(z, t) = \sum_k \alpha_k(t) e^{ikz} + c.c.$$

complex amplitude for mode e^{ikz}



Define

$$X(t) = \text{Re}[\alpha_k(t)] = \frac{1}{2} [\alpha_k(t) + \alpha_k^*(t)] = Q(t)$$

$$Y(t) = \text{Im}[\alpha_k(t)] = \frac{1}{2i} [\alpha_k(t) - \alpha_k^*(t)] = P(t)$$

Quantization: $\alpha \rightarrow \hat{a}, \alpha^* \rightarrow \hat{a}^\dagger$

$$\left. \begin{aligned} \hat{X}(t) &= \frac{1}{2} [\hat{a}_k(t) + \hat{a}_k^\dagger(t)] = \hat{Q}(t) \\ \hat{Y}(t) &= \frac{1}{2i} [\hat{a}_k(t) - \hat{a}_k^\dagger(t)] = \hat{P}(t) \end{aligned} \right\} [\hat{X}(t), \hat{Y}(t)] = i/2$$

$$\begin{aligned} \hat{E}(z, t) &= \mathcal{E}_k (\hat{X}(t) + i\hat{Y}(t)) e^{ikz} + \text{H.C.} \\ &= \mathcal{E}_k [\hat{X}(t) \cos(kz) - \hat{Y}(t) \sin(kz)] \end{aligned}$$

– same info, easier to work with –

Quantum States of the Field in Mode k

Number States (Fock states)

$$\hat{a}^\dagger \hat{a} |n\rangle = n |n\rangle$$



$$\langle n | \hat{X} |n\rangle = \langle n | \hat{Y} |n\rangle = 0$$

$$\langle n | \hat{X}^2 |n\rangle = \langle n | \hat{Y}^2 |n\rangle = \frac{1}{2} (n + 1/2)$$



$$\Delta X \Delta Y = \frac{1}{2} (n + 1/2)$$

- HIGHLY non-classical, $\langle \hat{E} \rangle = 0$
- VERY hard to make for large n

Quantum States of the Quantized Field

Quantum States of the Field in Mode k

Number States (Fock states)

$$\hat{a}^\dagger \hat{a} |n\rangle = n |n\rangle$$



$$\langle n | \hat{X} |n\rangle = \langle n | \hat{Y} |n\rangle = 0$$

$$\langle n | \hat{X}^2 |n\rangle = \langle n | \hat{Y}^2 |n\rangle = \frac{1}{2} (n + \frac{1}{2})$$



$$\Delta X \Delta Y = \frac{1}{2} (n + \frac{1}{2})$$

- HIGHLY non-classical, $\langle \hat{E} \rangle = 0$
- VERY hard to make for large n

Coherent States (Quasi-classical states)

- Closest approximation to classical field
- See Cohen-Tannoudj, complement G_V

Definition: $|\psi\rangle$ is coherent (quasiclassical) iff

$$\langle \hat{X}(t) \rangle = \langle \psi | \hat{X}(t) | \psi \rangle = X(t), \quad \langle \hat{Y}(t) \rangle = Y(t)$$

$$\langle \hat{H}(t) \rangle = \hbar \omega (|\alpha(t)|^2 + \frac{1}{2})$$

noting that

$$\hat{X}(t) \propto \hat{a}(t) = \hat{a}(0) e^{-i\omega t}$$

$$\hat{Y}(t) \propto \hat{a}^\dagger(t) = \hat{a}^\dagger(0) e^{i\omega t}$$



equivalently

Definition: $|\psi\rangle$ is coherent (quasiclassical) iff

$$(1) \quad \langle \hat{a}(0) \rangle = \langle \psi | \hat{a}(0) | \psi \rangle = \alpha(0)$$

$$(2) \quad \langle \hat{a}^\dagger(0) \hat{a}(0) \rangle = \alpha(0)^* \alpha(0)$$

Quantum States of the Quantized Field

Coherent States (Quasi-classical states)

- Closest approximation to classical field
- See Cohen-Tannoudj, complement G_v

Definition: $|\psi\rangle$ is coherent (quasiclassical) iff

$$\langle \hat{X}(t) \rangle = \langle \psi | \hat{X}(t) | \psi \rangle = X(t), \quad \langle \hat{Y}(t) \rangle = Y(t)$$

$$\langle \hat{H}(t) \rangle = \hbar\omega(|\alpha(t)|^2 + 1/2)$$

noting that

$$\hat{X}(t) \propto \hat{a}(t) = \hat{a}(0) e^{-i\omega t}$$

$$\hat{Y}(t) \propto \hat{a}^\dagger(t) = \hat{a}^\dagger(0) e^{i\omega t}$$

equivalently

Definition: $|\psi\rangle$ is coherent (quasiclassical) iff

$$(1) \quad \langle \hat{a}(0) \rangle = \langle \psi | \hat{a}(0) | \psi \rangle = \alpha(0)$$

$$(2) \quad \langle \hat{a}^\dagger(0) \hat{a}(0) \rangle = \alpha(0)^* \alpha(0)$$

Cohen-Tannoudji, Lecture Notes

equivalently

Definition: a state $|\alpha\rangle$ is coherent iff

$$\hat{a} |\alpha\rangle = \alpha |\alpha\rangle$$

Finally, one can show

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle$$


Physical properties

$$\langle \hat{X}(t) \rangle = \text{Re}[\alpha(0) e^{-i\omega t}]$$

$$\langle \hat{Y}(t) \rangle = \text{Im}[\alpha(0) e^{-i\omega t}]$$

$$\Delta X(t) = \Delta Y(t) = 1/2$$

$$\Delta X \Delta Y = 1/4$$



Quantum States of the Quantized Field

Cohen-Tannoudji, Lecture Notes



equivalently

Definition: a state $|\alpha\rangle$ is coherent iff

$$\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$$

Finally, one can show

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle$$


Physical properties

$$\langle \hat{X}(t) \rangle = \text{Re}[\alpha(0) e^{-i\omega t}]$$

$$\langle \hat{Y}(t) \rangle = \text{Im}[\alpha(0) e^{-i\omega t}]$$

$$\Delta X(t) = \Delta Y(t) = 1/2$$

$$\Delta X \Delta Y = 1/4$$



Photon statistics

Measure $\hat{N} \rightarrow$ $\left\{ \begin{array}{l} \text{outcomes } n \\ \mathcal{P}(n) = \langle \alpha | n \rangle \langle n | \alpha \rangle = \frac{|\alpha|^{2n}}{n!} e^{-|\alpha|^2} \end{array} \right.$



Poisson distribution w/ $\left\{ \begin{array}{l} \text{mean } \bar{n} = |\alpha|^2 \\ \text{variance } \Delta n^2 = |\alpha|^2 \end{array} \right.$



$$\Delta n = \sqrt{\bar{n}} \quad \text{-- Shot Noise}$$

Quantum States of the Quantized Field

Photon statistics

Measure \hat{N} \rightarrow { outcomes n


$$P(n) = \langle \alpha | n \times n | \alpha \rangle = \frac{|\alpha|^{2n}}{n!} e^{-|\alpha|^2}$$

Poisson distribution w/ {

- mean $\bar{n} = |\alpha|^2$
- variance $\Delta n^2 = |\alpha|^2$

$$\Delta n = \sqrt{\bar{n}} \quad \text{-- Shot Noise}$$

More about Coherent States



Coherent States as translated Vacuum States?

Generating Coherent States from the Vacuum

Definition: $\hat{D}(\alpha) = e^{\alpha \hat{a}^\dagger - \alpha^* \hat{a}}$

Unitary, equals translation


Glaubers formula (from BCH formula)

$$e^{\hat{A} + \hat{B}} = e^{\hat{A}} e^{\hat{B}} e^{\frac{1}{2}[\hat{A}, \hat{B}]}$$

for $[\hat{A}, [\hat{A}, \hat{B}]] = [\hat{B}, [\hat{A}, \hat{B}]] = 0$

Quantum States of the Quantized Field

More about Coherent States



Coherent States
as translated
Vacuum States?

Generating Coherent States from the Vacuum

Definition: $\hat{D}(\alpha) = e^{\alpha \hat{a}^\dagger - \alpha^* \hat{a}}$

Unitary, equals translation

Glauber's formula (from BCH formula)

$$e^{\hat{A} + \hat{B}} = e^{\hat{A}} e^{\hat{B}} e^{\frac{i}{2} [\hat{A}, \hat{B}]}$$

for $[\hat{A}, [\hat{A}, \hat{B}]] = [\hat{B}, [\hat{A}, \hat{B}]] = 0$

Apply to

$$[\alpha \hat{a}^\dagger, -\alpha^* \hat{a}] = \alpha^* \alpha$$

\uparrow \uparrow \uparrow
 \hat{A} \hat{B} $[\hat{A}, \hat{B}]$

$$\hat{D}(\alpha) = e^{-|\alpha|^2/2} e^{\alpha \hat{a}^\dagger} e^{-\alpha \hat{a}}$$

Remember: $\hat{a}|0\rangle = 0$

$$e^{-\alpha \hat{a}} |0\rangle = \sum_n \frac{(-\alpha \hat{a})^n}{n!} |0\rangle = |0\rangle$$

$$\begin{aligned} \hat{D}(\alpha)|0\rangle &= e^{-|\alpha|^2/2} e^{\alpha \hat{a}^\dagger} |0\rangle \\ &= e^{-|\alpha|^2} \sum_n \frac{(\alpha \hat{a}^\dagger)^n}{n!} |0\rangle \\ &= e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle = |\alpha\rangle \end{aligned}$$

$$\hat{D}(\alpha)|0\rangle = |\alpha\rangle$$

Quantum States of the Quantized Field

Apply to

$$[\alpha \hat{a}^\dagger, -\alpha^* \hat{a}] = \alpha^* \alpha$$

\uparrow \uparrow \uparrow
 \hat{A} \hat{B} $[\hat{A}, \hat{B}]$



$$\hat{D}(\alpha) = e^{-|\alpha|^2/2} e^{\alpha \hat{a}^\dagger} e^{-\alpha \hat{a}}$$

Remember:

$$\hat{a}|0\rangle = 0 \rightarrow$$

$$e^{-\alpha^* \hat{a}} |0\rangle = \sum_n \frac{(-\alpha^* \hat{a})^n}{n!} |0\rangle = |0\rangle$$



$$\begin{aligned} \hat{D}(\alpha)|0\rangle &= e^{-|\alpha|^2/2} e^{\alpha \hat{a}^\dagger} |0\rangle \\ &= e^{-|\alpha|^2} \sum_n \frac{(\alpha \hat{a}^\dagger)^n}{n!} |0\rangle \\ &= e^{-|\alpha|^2/2} \sum_n \frac{\alpha^n}{\sqrt{n!}} |n\rangle = |\alpha\rangle \end{aligned}$$



$$\hat{D}(\alpha)|0\rangle = |\alpha\rangle$$

OK – $\hat{D}(\alpha)$ generates $|\alpha\rangle$ from the vacuum!

Rewrite:

$$\begin{aligned} \alpha \hat{a}^\dagger - \alpha^* \hat{a} &= (\alpha - \alpha^*) \hat{X} + i(\alpha + \alpha^*) \hat{Y} \\ &= i2Y\hat{X} + i2X\hat{Y} \end{aligned}$$

where $X = \langle \alpha | \hat{X} | \alpha \rangle$, $Y = \langle \alpha | \hat{Y} | \alpha \rangle$

Glauber's formula again:

$$\hat{D}(\alpha) = e^{i2Y\hat{X} + i2X\hat{Y}} = e^{-XY/4} e^{i2Y\hat{X}} e^{i2X\hat{Y}}$$

Recall: $\hat{S}(q) = e^{-iq\hat{P}/\hbar} \rightarrow$ translation by q

$\hat{S}(p) = e^{-ip\hat{Q}/\hbar} \rightarrow$ translation by p

where

$$q = q_0 X, \quad p = p_0 Y$$

$$\hat{q} = q_0 \hat{X}, \quad \hat{p} = p_0 \hat{Y}$$

$$\& \quad X_0 p_0 = 2\hbar$$

Quantum States of the Quantized Field

OK – $\hat{D}(\alpha)$ generates $|\alpha\rangle$ from the vacuum!

Rewrite:

$$\alpha \hat{a}^\dagger - \alpha^* \hat{a} = (\alpha - \alpha^*) \hat{X} + i(\alpha + \alpha^*) \hat{Y} \\ = i2Y\hat{X} + i2X\hat{Y}$$

where $X = \langle \alpha | \hat{X} | \alpha \rangle$, $Y = \langle \alpha | \hat{Y} | \alpha \rangle$

Glauber's formula again:

$$\hat{D}(\alpha) = e^{i2Y\hat{X} + i2X\hat{Y}} = e^{-XY/4} e^{i2Y\hat{X}} e^{i2X\hat{Y}}$$

Recall: $\hat{S}(q) = e^{-iq\hat{P}/\hbar}$ → translation by q


Recall: $\hat{S}(p) = e^{-ip\hat{Q}/\hbar}$ → translation by p

where $q = q_0 X$, $p = p_0 Y$ & $X_0 p_0 = 2\hbar$
 $\hat{q} = q_0 \hat{X}$, $\hat{p} = p_0 \hat{Y}$

This gives us

$$\hat{S}(q) = \hat{S}(X) = e^{i2X\hat{Y}}, \quad \hat{S}(p) = \hat{S}(Y) = e^{i2Y\hat{X}}$$

$\hat{D}(\alpha)$ translates
along X then Y




**Discussion –
How to do this?**

Quantum States of the Quantized Field

This gives us

$$\hat{S}(q) = \hat{S}(x) = e^{i2x\hat{Y}}, \quad \hat{S}(p) = \hat{S}(y) = e^{i2y\hat{X}}$$

$\hat{D}(\alpha)$ translates
along X then Y



**Discussion –
How to do this?**

Coherent States from Classical Dipole Radiation

Classical Dipole $d(t) = d_0 \cos(\omega t)$ @ $t=0$

Quantized Field $\hat{E}(z) = \epsilon_{\mathbf{k}} (\hat{a} + \hat{a}^\dagger)$

Dipole-Field Interaction

$$\hat{H} = \hbar\omega_{\mathbf{k}} (\hat{a}^\dagger \hat{a} + 1/2) + \hbar\lambda(t) (\hat{a} + \hat{a}^\dagger)$$

$$\lambda(t) = -\frac{d(t)\epsilon_{\mathbf{k}}}{\hbar} = \lambda_0 \cos(\omega t)$$

**Homework Problem
(voluntary)**

$$\alpha(T) = -i \frac{\lambda_0}{2} e^{-i(\omega - \omega_{\mathbf{k}})T/2} \frac{\sin[(\omega - \omega_{\mathbf{k}})T/2]}{(\omega - \omega_{\mathbf{k}})/2}$$

Quantum States of the Quantized Field

Coherent States from Classical Dipole Radiation

Classical Dipole $d(t) = d_0 \cos(\omega t)$ @ $t=0$

Quantized Field $\hat{E}(z) = \epsilon_k (\hat{a} + \hat{a}^\dagger)$

Dipole-Field Interaction


$$\hat{H} = \hbar\omega (\hat{a}^\dagger \hat{a} + 1/2) + \hbar\lambda(t) (\hat{a} + \hat{a}^\dagger)$$

$$\lambda(t) = -\frac{d(t)\epsilon_k}{\hbar} = \lambda_0 \cos(\omega t)$$



Homework Problem
(voluntary)

$$\alpha(T) = -i \frac{\lambda_0}{2} e^{-i(\omega - \omega_k)T/2} \frac{\sin[(\omega - \omega_k)T/2]}{(\omega - \omega_k)/2}$$



Recall from Semi-Classical Laser Theory

$$\langle \hat{j}(t) \rangle \text{ drives } \hat{E}(t)$$

classical dipole
+ quantum
fluctuations

coherent state
+ quantum
fluctuations

Drive from $0 < t < T$ →

$$\alpha(t) = \alpha(T) e^{-i\omega_k(t-T)}$$

