Begin
02-20-2023

Emission and Absorption – Population Rate Equations

So far we have derived a set of Eqs. of Motion for the elements of the Density Matrix:

$$\dot{\rho}_{11} = -\Gamma_1 \rho_{11} + A_{21}\rho_{22} - \frac{i}{2}(\chi\rho_{12} - \chi^*\rho_{21})$$

$$\dot{\rho}_{22} = -\Gamma_2 \rho_{22} - A_{21}\rho_{22} + \frac{i}{2}(\chi\rho_{12} - \chi^*\rho_{21})$$

$$\dot{\rho}_{12} = (i\Delta - \beta)\rho_{12} + \frac{i\chi^*}{2}(\rho_{22} - \rho_{11}) = \dot{\rho}_{21}^*$$

where $\beta = \frac{1}{\tau} + \frac{A_{21}}{2} + \frac{\Gamma_1 + \Gamma_2}{2}$

(*) These eqs. are difficult to solve in the general case. See, e. g., Allen & Eberly for discussion of some special cases and a reference to original work by Torrey et al.

(*) For ≥ 3 levels the Density Matrix Equations get very cumbersome and it is desirable to simplify the description when possible.

(*) One such simplification takes the form of Rate Equations for the populations only.

Steady State Solutions: (requires $\Gamma_1 = \Gamma_2 = 0$)

Let $\dot{\rho}_{12} = 0$ ⇒

$$\begin{cases} \rho_{12} = \dfrac{i\chi^*/2}{\beta - i\Delta}(\rho_{22} - \rho_{11}) \\ \rho_{21} = \dfrac{-i\chi/2}{\beta + i\Delta}(\rho_{22} - \rho_{11}) \end{cases}$$

$$\chi\rho_{12} - \chi^*\rho_{21} = \frac{i|\chi|^2\beta}{\Delta^2 + \beta^2}(\rho_{22} - \rho_{11})$$

Plug into eqs for populations to get

$$\dot{\rho}_{11} = A_{21}\rho_{22} + \frac{|\chi|^2\beta/2}{\Delta^2 + \beta^2}(\rho_{22} - \rho_{11}) = 0$$

$$\dot{\rho}_{22} = -A_{21}\rho_{22} - \frac{|\chi|^2\beta/2}{\Delta^2 + \beta^2}(\rho_{22} - \rho_{11}) = 0$$

From these eqs. we can find <u>steady state values</u> for the populations and coherences in terms of $\chi, \Delta, A_{21}, \beta$ when (and only when) $\dot{\rho}_{11} = \dot{\rho}_{22} = 0$

Emission and Absorption – Population Rate Equations

Steady State Solutions: (requires $\Gamma_1 = \Gamma_2 = 0$)

Let $\dot{\rho}_{12} = 0$ ⇒

$$\begin{cases} \rho_{12} = \dfrac{i\chi^*/2}{\beta - i\Delta}(\rho_{22} - \rho_{11}) \\ \rho_{21} = \dfrac{-i\chi/2}{\beta + i\Delta}(\rho_{22} - \rho_{11}) \end{cases}$$

$$\chi\rho_{12} - \chi^*\rho_{21} = \frac{i|\chi|^2\beta}{\Delta^2 + \beta^2}(\rho_{22} - \rho_{11})$$

Plug into eqs for populations to get

$$\dot{\rho}_{11} = A_{21}\rho_{22} + \frac{|\chi|^2\beta/2}{\Delta^2 + \beta^2}(\rho_{22} - \rho_{11}) = 0$$

$$\dot{\rho}_{22} = -A_{21}\rho_{22} - \frac{|\chi|^2\beta/2}{\Delta^2 + \beta^2}(\rho_{22} - \rho_{11}) = 0$$

From these eqs. we can find steady state values for the populations and coherences in terms of $\chi, \Delta, A_{21}, \beta$ when (and only when) $\dot{\rho}_{11} = \dot{\rho}_{22} = 0$

Note: The terms remaining after elimination of ρ_{12}, ρ_{21} are commonly identified with induced or stimulated processes. They are proportional to $|\chi|^2$, $|E_0|^2$ and thus the intensity of the light field.

Def: Absorption Rate = Stimulated Emission Rate

$$R_{12} = \frac{|\chi|^2\beta/2}{\Delta^2 + \beta^2} = \frac{|\vec{\mu}_{12}\cdot\vec{\mathcal{E}}E_0/\hbar|^2\beta/2}{(\omega_{21} - \omega)^2 + \beta^2}$$

Schematic:



Emission and Absorption – Population Rate Equations

Note: The terms remaining after elimination of ρ_{12}, ρ_{21} are commonly identified with induced or stimulated processes. They are proportional to $|X|^2, |E_0|^2$ and thus the intensity of the light field.

Def: Absorption Rate = Stimulated Emission Rate

$$R_{12} = \frac{|X|^2 \beta/2}{\Delta^2 + \beta^2} = \frac{|\vec{\mu}_{12} \cdot \vec{\mathcal{E}} E_0/\hbar|^2 \beta/2}{(\omega_{21} - \omega)^2 + \beta^2}$$

Schematic:



Elastic Collision Broadening

In hot and dense gases the dominant source of relaxation is often elastic collisions between atoms

Let $\beta \gg \Gamma_1, \Gamma_2, A_{21}$ ⇒ ρ_{12} reaches steady state much faster than ρ_{11}, ρ_{22}

We can solve the eq. for $\dot{\rho}_{12}$ assuming it is in steady state for given values of ρ_{11}, ρ_{22}

This yields Rate Equations for the populations only, valid in the collision broadened regime

$$\dot{\rho}_{11} = -\Gamma_1 \rho_{11} + A_{21} \rho_{22} + R_{12}(\rho_{22} - \rho_{11}) \neq 0$$

$$\dot{\rho}_{22} = -\Gamma_2 \rho_{22} - A_{21} \rho_{22} - R_{12}(\rho_{22} - \rho_{11}) \neq 0$$

(*) This is another example of adiabatic elimination of a fast variable (the coherence), leaving us with simpler equations for the slower variables.

(*) From these we can find the transient behavior of the coherences ρ_{11}, ρ_{22}

Emission and Absorption – Population Rate Equations

Elastic Collision Broadening

In hot and dense gases the dominant source of relaxation is often elastic collisions between atoms

Let $\beta \gg \Gamma_1, \Gamma_2, A_{21}$ ⇒ σ_{12} reaches steady state much faster than σ_{11}, σ_{22}

We can solve the eq. for $\dot{\sigma}_{12}$ assuming it is in steady state for given values of σ_{11}, σ_{22}

This yields Rate Equations for the populations only, valid in the collision broadened regime

$$\dot{\sigma}_{11} = -\Gamma_1 \sigma_{11} + A_{21}\sigma_{22} + R_{12}(\sigma_{22} - \sigma_{11}) \neq 0$$

$$\dot{\sigma}_{22} = -\Gamma_2 \sigma_{22} - A_{21}\sigma_{22} - R_{12}(\sigma_{22} - \sigma_{11}) \neq 0$$

(*) This is another example of adiabatic elimination of a fast variable (the coherence), leaving us with simpler equations for the slower variables.

(*) From these we can find the transient behavior of the coherences σ_{11}, σ_{22}

Note: When collisions are very frequent the dipole $\langle \hat{\mu} \rangle$ is oriented at random relative to the driving field. In that case

$$\langle |\vec{\mu}_{12} \cdot \vec{\mathcal{E}} E_0|^2 \rangle_{angles} = \frac{1}{3} \mu_{12}^2 |E_0|^2 \Rightarrow$$

$$R_{12} = \frac{\langle |\vec{\mu}_{12} \cdot \vec{\mathcal{E}} E_0/\hbar|^2 \rangle_{angles}\, \beta/2}{\Delta^2 + \beta^2} = \frac{1}{3} \frac{|\chi|^2 \beta/2}{\Delta^2 + \beta^2}$$

Photon Flux and Cross Section

Let $R_{12} \equiv \sigma(\Delta)\phi$ where $\hbar\omega\phi = \frac{1}{2} c\varepsilon_0 |E_0|^2$

"photon flux" — intensity

This allows us to recast the Rate Eqs

$$\dot{\sigma}_{11} = -\Gamma_1 \sigma_{11} + A_{21}\sigma_{22} + \sigma(\Delta)\phi(\sigma_{22} - \sigma_{11})$$

$$\dot{\sigma}_{22} = -\Gamma_2 \sigma_{22} - A_{21}\sigma_{22} - \sigma(\Delta)\phi(\sigma_{22} - \sigma_{11})$$

Emission and Absorption – Population Rate Equations

Note: When collisions are very frequent the dipole $\langle \hat{\vec{\mu}} \rangle$ is oriented at random relative to the driving field. In that case

$$\langle |\vec{\mu}_{12} \cdot \vec{\mathcal{E}} E_0|^2 \rangle_{angles} = \frac{1}{3} \mu_{12}^2 |E_0|$$

$$R_{12} = \frac{\langle |\vec{\mu}_{12} \cdot \vec{\mathcal{E}} E_0/\hbar|^2 \rangle_{angles} \beta/2}{\Delta^2 + \beta^2} = \frac{1}{3} \frac{|\chi|^2 \beta/2}{\Delta^2 + \beta^2}$$

Photon Flux and Cross Section

Let $R_{12} \equiv \sigma(\Delta)\phi$ where $\hbar\omega\phi = \underbrace{\frac{1}{2} c \varepsilon_0 |E_0|^2}_{\text{intensity}}$

"photon flux"

This allows us to recast the Rate Eqs

$$\dot{\varrho}_{11} = -\Gamma_1 \varrho_{11} + A_{21} \varrho_{22} + \sigma(\Delta)\phi(\varrho_{22} - \varrho_{11})$$

$$\dot{\varrho}_{22} = -\Gamma_2 \varrho_{22} - A_{21} \varrho_{22} - \sigma(\Delta)\phi(\varrho_{22} - \varrho_{11})$$

We see that per atom, per unit time

# of absorption events	=	$\sigma(\Delta)\phi\varrho_{11}$
# of stim. emission events	=	$\sigma(\Delta)\phi\varrho_{22}$

Note: Given N atoms, the total # of events are $N\varrho_{11}$, $N\varrho_{22}$. This is useful when we care about the total power in the light field, e. g., in the context of laser theory

Solution of the Rate Equations

Let $\Gamma_1 = \Gamma_2 = 0$ and plug in $\varrho_{11} = 1 - \varrho_{22}$

$$\dot{\varrho}_{22} = -A_{21}\varrho_{22} - \sigma(\Delta)\phi(2\varrho_{22} - 1)$$

$$= -\underbrace{(A_{21} + 2\sigma(\Delta)\phi)}_{\gamma} \varrho_{22} + \sigma(\Delta)\phi$$

This solution is a damped approach to Steady State!

Emission and Absorption – Population Rate Equations

We see that per atom, per unit time

# of absorption events	=	$\sigma(\Delta)\phi\,\rho_{11}$
# of stim. emission events	=	$\sigma(\Delta)\phi\,\rho_{22}$

Note: Given N atoms, the total # of events are $N\rho_{11}$, $N\rho_{22}$. This is useful when we care about the total power in the light field, e. g., in the context of laser theory

Solution of the Rate Equations

Let $\Gamma_1 = \Gamma_2 = 0$ and plug in $\rho_{11} = 1 - \rho_{22}$

$$\dot{\rho}_{22} = -A_{21}\rho_{22} - \sigma(\Delta)\phi\,(2\rho_{22} - 1)$$
$$= -\underbrace{(A_{21} + 2\sigma(\Delta)\phi)}_{\gamma}\rho_{22} + \sigma(\Delta)\phi$$

The solution is a damped approach to Steady State!

$$\rho_{22}(t) = [\rho_{22}(0) - \rho_{22}(\infty)]e^{-\gamma t} + \rho_{22}(\infty)$$

where

$$\gamma = (A_{21} + 2\sigma(\Delta)\phi)\,,\quad \rho_{22}(\infty) = \frac{\sigma(\Delta)\phi}{A_{21} + 2\sigma(\Delta)\phi}$$



(*) **This transient behavior is valid in the collision broadened regime.**

(*) Without collisions the transient regime Is one of damped Rabi oscillations.

(*) The steady state value $\rho_{22}(\infty)$ is good regardless

Emission and Absorption – Population Rate Equations

$$\rho_{22}(t) = [\rho_{22}(0) - \rho_{22}(\infty)]e^{-\gamma t} + \rho_{22}(\infty)$$

where

$$\gamma = (A_{21} + 2\sigma(\Delta)\phi)\ ,\quad \rho_{22}(\infty) = \frac{\sigma(\Delta)\phi}{A_{21} + 2\sigma(\Delta)\phi}$$



(*) This <u>transient</u> behavior is valid in the collision broadened regime.

(*) Without collisions the transient regime Is one of damped Rabi oscillations.

(*) The <u>steady state</u> value $\rho_{22}(\infty)$ is good regardless

Numerical simulation of Density Matrix Eqs (Optical Bloch Picture).

Figure from Milloni & Eberly



Figure 6.6 Numerical solutions of the v, w equations (6.5.21) for a range of collisional damping rates. Note scale changes.

Emission and Absorption – Population Rate Equations

$$\rho_{22}(t) = [\rho_{22}(0) - \rho_{22}(\infty)]\, e^{-\gamma t} + \rho_{22}(\infty)$$

where

$$\gamma = (A_{21} + 2\sigma(\Delta)\phi)\ ,\quad \rho_{22}(\infty) = \frac{\sigma(\Delta)\phi}{A_{21} + 2\sigma(\Delta)\phi}$$



(*) This <u>transient</u> behavior is valid in the collision broadened regime.

(*) Without collisions the transient regime Is one of damped Rabi oscillations.

(*) The <u>steady state</u> value $\rho_{22}(\infty)$ is good regardless

Limiting cases:

$\sigma(\Delta)\phi = 0 \Rightarrow \rho_{22}(\infty) = 0$

$\sigma(\Delta)\phi \ll A_{21} \Rightarrow \rho_{22}(\infty) = \dfrac{\sigma(\Delta)\phi}{A_{21}}$

$\sigma(\Delta)\phi \gg A_{21} \Rightarrow \rho_{22}(\infty) = 1/2$ ← Saturation!

Rewrite $\rho_{22}(\infty)$ using $R_{12} = \sigma(\Delta)\phi = \dfrac{|X|^2\beta/2}{\Delta^2 + \beta^2}$

$$\Rightarrow \rho_{22}(\infty) = \frac{|X|^2\beta/2A_{21}}{\Delta^2 + \beta^2 + |X|^2\beta/A_{21}}$$

Plot $\rho_{22}(\infty)$ vs Δ:



HWHM line width:

$$\Delta_{1/2} = \sqrt{\beta^2 + |X|^2\beta/A_{21}}$$

$$= \beta\sqrt{1 + \frac{2\sigma(0)\phi}{A_{21}}}$$

$$\left(\text{used } \sigma(0)\phi = \frac{|X|^2}{2\beta}\right)$$

Emission and Absorption – Population Rate Equations

Limiting cases:

$\sigma(\Delta)\phi = 0 \Rightarrow \rho_{22}(\infty) = 0$

$\sigma(\Delta)\phi \ll A_{21} \Rightarrow \rho_{22}(\infty) = \dfrac{\sigma(\Delta)\phi}{A_{21}}$

$\sigma(\Delta)\phi \gg A_{21} \Rightarrow \rho_{22}(\infty) = 1/2$ ← Saturation!

Rewrite $\rho_{22}(\infty)$ using $R_{12} = \sigma(\Delta)\phi = \dfrac{|X|^2\beta/2}{\Delta^2+\beta^2}$

$$\Rightarrow \rho_{22}(\infty) = \frac{|X|^2\beta/2\,A_{21}}{\Delta^2+\beta^2+|X|^2\beta/A_{21}}$$

Plot $\rho_{22}(\infty)$ vs Δ:



HWHM line width:

$$\Delta_{1/2} = \sqrt{\beta^2+|X|^2\beta/A_{21}}$$

$$= \beta\sqrt{1+\frac{2\sigma(0)\phi}{A_{21}}}$$

$\left(\text{used } \sigma(0)\phi = \dfrac{|X|^2}{2\beta}\right)$

Power Broadening: Rewrite

$$\Delta_{1/2} = \beta\sqrt{1+\phi/\phi_{SAT}} = \beta\sqrt{1+I/I_{SAT}}$$

where

$$\phi_{SAT} \equiv \frac{A_{21}}{2\sigma(0)}, \quad I_{SAT} \equiv \frac{\hbar\omega A_{21}}{2\sigma(0)}$$

β : natural linewidth

Power Broadening in molecular beam spectroscopy:



Keep $I \ll I_{SAT}$ for best spectroscopic resolution

Emission and Absorption – Population Rate Equations

Begin
02-22-2023

Power Broadening: Rewrite

$$\Delta_{1/2} = \beta\sqrt{1+\phi/\phi_{SAT}} = \beta\sqrt{1+I/I_{SAT}}$$

where

$$\phi_{SAT} \equiv \frac{A_{21}}{2\sigma(0)} \quad , \quad I_{SAT} \equiv \frac{\hbar\omega A_{21}}{2\sigma(0)}$$

β : natural linewidth

Power Broadening in molecular beam spectroscopy:



Keep $I \ll I_{SAT}$ for best spectroscopic resolution

End
02-20-2023

More about the Photon Scattering Cross Section

By Definition

$$R_{12} = \frac{|X_1|^2\beta/2}{\Delta^2+\beta^2} = \sigma(\Delta)\phi = \sigma(\Delta)\frac{1/2\,c\varepsilon_0|E_0|^2}{\hbar\omega}$$

where $|X|^2 = |\vec{\mu}_{12}\cdot\vec{\varepsilon}E_0/\hbar|^2 = f\frac{|\vec{\mu}_{12}|^2|E_0|^2}{\hbar^2}$

and Collision free → $1 \geq f \geq 1/3$ ← Collision broadened

This gives us

$$\sigma(\Delta) = f\frac{\omega\mu_{12}^2}{\hbar c\varepsilon_0\beta}\frac{\beta^2}{\Delta^2+\beta^2} \equiv \sigma(0)\frac{\beta^2}{\Delta^2+\beta^2}$$

Let $\beta = A_{21}/2$, $\Gamma_1 = \Gamma_2 = 0$ (collision free)

$$\sigma(0) = f\frac{2\omega\mu_{12}^2}{\hbar c\varepsilon_0 A_{21}} = f\frac{4\pi}{\hbar\varepsilon_0\lambda}\frac{\mu_{21}^2}{A_{21}} \quad *$$

The connection between A_{21} and μ_{21}^2 is intuitive, derived rigorously during the QED part of OPTI 544

Emission and Absorption – Population Rate Equations

Begin
02-22-2023

More about the Photon Scattering Cross Section

By Definition

$$R_{12} = \frac{|X_1|^2 \beta/2}{\Delta^2 + \beta^2} = \sigma(\Delta)\phi = \sigma(\Delta)\frac{1/2\, c\varepsilon_0 |E_0|^2}{\hbar\omega}$$

where $|X|^2 = |\vec{p}_{12}\cdot\vec{\mathcal{E}} E_0/\hbar|^2 = f\frac{|\vec{p}_{12}|^2 |E_0|^2}{\hbar^2}$

and Collision free → $1 \geq f \geq 1/3$ ← Collision broadened

This gives us

$$\sigma(\Delta) = f\frac{\omega p_{12}^2}{\hbar c\varepsilon_0 \beta}\frac{\beta^2}{\Delta^2+\beta^2} \equiv \sigma(0)\frac{\beta^2}{\Delta^2+\beta^2}$$

Let $\beta = A_{21}/2$, $\Gamma_1 = \Gamma_2 = 0$ (collision free)

$$\sigma(0) = f\frac{2\omega p_{12}^2}{\hbar c\varepsilon_0 A_{21}} = f\frac{4\pi}{\hbar\varepsilon_0\lambda}\frac{p_{21}^2}{A_{21}} \quad *$$

The connection between A_{21} and p_{21}^2 is intuitive, derived rigorously during the QED part of OPTI 544

End
02-20-2023

Here we simply note the result :

$$A_{21} = \frac{p_{12}^2 \omega^3}{3\pi\varepsilon_0\hbar c^3}$$

Substituting in * we get

$$\sigma(0) = f\frac{2\omega}{\hbar c\varepsilon_0}\frac{3\pi\varepsilon_0\hbar c^3}{\omega^3} = f\frac{3\lambda^2}{2\pi}$$

$$\frac{3\lambda^2}{2\pi} \geq \sigma(0) \geq \frac{\lambda^2}{2\pi}$$

Collision free, polarized light — Collision broadened or un-polarized light

– Remarkably simple result – easy to remember

Why ?

Emission and Absorption – Population Rate Equations

Here we simply note the result :

$$A_{21} = \frac{n_{12}^2 \omega^3}{3\pi\varepsilon_0 \hbar c^3}$$

Substituting in $*$ we get

$$\sigma(0) = f\frac{2\omega}{\hbar c \varepsilon_0}\frac{3\pi\varepsilon_0\hbar c^3}{\omega^3} = f\frac{3\lambda^2}{2\pi}$$

$$\frac{3\lambda^2}{2\pi} \geq \sigma(0) \geq \frac{\lambda^2}{2\pi}$$

Collision free, polarized light

Collision broadened or un-polarized light

– Remarkably simple result – easy to remember

Why ?

Rate Eq. Model of Absorption

Remember our Physical Picture of absorption/ stim. emission



The loss over distance dz is $d\phi = \sigma(\Delta)\phi N(\varrho_{22} - \varrho_{11})dz$

If $\phi \ll \phi_{sat}$ then $\varrho_{11} \simeq 1$ and $\varrho_{22} \simeq 0$ and we get

$$d\phi = -N\sigma(\Delta)\phi dz \Rightarrow \frac{d}{dz}\phi = -N\sigma(\Delta)\phi = -a\phi$$

Absorption:



$$\phi(z) = e^{-a(z-z_0)}\phi(z_0)$$

$$I(z) = e^{-a(z-z_0)}I(z_0)$$

Typical $\lambda = 500\,nm$, $a = N \times 10^{-9}\,cm^2 \frac{\beta^2}{\Delta^2+\beta^2}$

Solid: $N \sim 10^{21}/cm^3$ ⇒ totally opaque @ resonance

Gas: $N \sim 10^{8}/cm^3$ ⇒ transparent @ resonance

Emission and Absorption – Population Rate Equations

Rate Eq. Model of Absorption

Remember our Physical Picture of absorption/ stim. emission



The loss over distance dz is $d\phi = \sigma(\Delta)\phi N(\rho_{22}-\rho_{11})dz$

If $\phi \ll \phi_{sat}$ then $\rho_{11}\simeq 1$ and $\rho_{22}\simeq 0$ and we get

$d\phi = -N\sigma(\Delta)\phi dz \Rightarrow \frac{d}{dz}\phi = -N\sigma(\Delta)\phi = -a\phi$

Absorption:



$$\phi(z) = e^{-a(z-z_0)}\phi(z_0)$$
$$I(z) = e^{-a(z-z_0)}I(z_0)$$

Typical $\lambda = 500\,nm$, $a = N\times 10^{-9}\,cm^2\frac{\beta^2}{\Delta^2+\beta^2}$

Solid: $N\sim 10^{21}/cm^3$ ⇒ totally opaque @ resonance

Gas: $N\sim 10^{8}/cm^3$ ⇒ transparent @ resonance

Note: This is a low saturation result !

High flux ⇒ photon scattering per atom saturates

If $\phi \gg \phi_{sat}$ then $\rho_{22}(\infty) = 1/2 \Rightarrow d\phi = N\frac{A_{21}}{2}dz$

$$\phi(z) = \phi(z_0) - \frac{NA_{21}}{2}(z-z_0)$$

At very high flux the loss is relatively insignificant

$$\phi(z), \phi(z_0) \gg \frac{NA_{21}}{2}(z-z_0)$$

– This is referred to as <u>bleaching</u> or <u>hole burning</u>

Emission and Absorption – Population Rate Equations

Blackbody Radiation

This is a standard problem in Optical Physics, which we review only briefly in OPTI 544. See almost any textbook for details, including Milloni & Eberly.

Step (1): We define a Lineshape Function:

$$\sigma(\Delta) = \frac{\pi \eta_{12}^2 \omega_{21}}{2\varepsilon_0 \hbar c} \underbrace{\frac{\delta\nu_{21}/\pi}{(\nu_{21}-\nu)^2 + \delta\nu_{21}^2}}_{S(\nu)}, \quad \delta\nu_{21} = \beta/2\pi$$

$S(\nu)$ is sharply peaked compared to the energy density $\rho(\nu)$ of Blackbody Radiation.

⇨ we can approximate $S(\nu) = \delta(\nu_{21} - \nu)$

Step (2): Sub in the expression for transition rates and integrate over all frequencies

$$R_{12} = \int_0^\infty \sigma(\nu)\phi(\nu)\,d\nu = \frac{\pi \eta_{21}^2 \omega_{21}}{2\varepsilon_0 \hbar} \int_0^\infty S(\nu) \frac{\rho(\nu)}{\hbar\nu}\,d\nu$$

$$\Rightarrow \quad R_{12} = \boxed{\frac{\pi \eta_{12}^2}{2\varepsilon_0 \hbar^2}} \rho(\nu_{21}) = B\rho(\nu_{21})$$

← Einstein B coefficient

Step (3): 2-level atom in thermal equilibrium with Blackbody Radiation field

$$\frac{g_2}{g_1} = e^{-h\nu_{21}/k_B T}$$

$$(A_{21} + R_{21})\rho_2 = R_{12}\rho_1$$

(Detailed Balance)

$$\Rightarrow \quad \boxed{\rho(\nu_{21}) = \frac{A_{21}/B}{e^{h\nu_{21}/k_B T} - 1}}$$

Step (4): Use prior result for A_{21} to find A_{21}/B which is independent of η_{12}^2

$$\boxed{\rho(\nu) = \frac{8\pi h\nu^3/c^3}{e^{h\nu/k_B T} - 1}}$$

Energy Density of the Blackbody Radiation field

Step (5): We can extend this to find the

* Total electromagnetic energy density
* Total radiated intensity from a black body

Emission and Absorption – Population Rate Equations

Step (3): 2-level atom in thermal equilibrium with Blackbody Radiation field

$$\frac{g_2}{g_1} = e^{-h\nu_{21}/k_B T}$$

$$(A_{21} + R_{21})\rho_2 = R_{12}\rho_1$$

(Detailed Balance)

$$\Rightarrow \boxed{\rho(\nu_{21}) = \frac{A_{21}/B}{e^{h\nu_{21}/k_B T} - 1}}$$

Step (4): Use prior result for A_{21} to find A_{21}/B, which is independent of μ_{12}^2

$$\boxed{\rho(\nu) = \frac{8\pi h\nu^3/c^3}{e^{h\nu/k_B T} - 1}}$$

Energy Density of the Blackbody Radiation field

Step (5): We can extend this to find the

* Total electromagnetic energy density
* Total radiated intensity from a black body

Step (6): Relative importance of spontaneous and stimulated emission

We have $\frac{A_{21}}{R_{12}} = \frac{A_{21}}{B\rho(\nu)} = e^{h\nu/k_B T} - 1 \gg 1$

for $T <$ several $\times$ 1000K

Take the surface of the Sun, $T = 5800K$

$$\boxed{\frac{A_{21}}{R_{12}} \sim 140 \;@\; \lambda_{max} = 500\,nm}$$

End 02-22-2023