

**OPTI 544: Problem Set 7**  
**Posted April 8, Due April 15**

**Electronic Submission Only, by email to Jon Pajaud ( jpajaud@email.arizona.edu )**

**I**

Consider the  $x$ -polarized electromagnetic field in a cylindrical cavity of length  $L$  and cross sectional area  $A$ . Using the standard normal mode expansion of  $E_x(z,t)$  and  $B_y(z,t)$  from class, write down expressions for the Hamiltonian and the Lagrangian in terms of the generalized coordinates  $q_j(t)$  and their time derivatives.

- (a) Use the Lagrange equations of motion to derive a second order differential equation for the  $q_j(t)$ 's.
- (b) Substitute the normal mode expansion of  $E_x(z,t)$  in the wave equation and derive a second order differential equation for the  $q_j(t)$ 's. Compare to the result in (a) above.

**II**

Consider in the following a 4-port beamsplitter with  $t = 1/\sqrt{2}$  and  $r = i/\sqrt{2}$ .

- (a) Let the input state be  $|\Psi_{in}\rangle = (\sqrt{1-\varepsilon}|1\rangle_1 + \sqrt{\varepsilon}|2\rangle_1)(\sqrt{1-\varepsilon}|1\rangle_2 + \sqrt{\varepsilon}|2\rangle_2)$ .
  - i. e. the wavepackets entering each port are mostly one-photon states but contain a small admixture of two-photon states. Find the output state  $|\Psi_{out}\rangle$ .

We use photomultiplier type detectors to measure the outputs from the beamsplitter. These detectors will click once when struck by a pulse of one or more photons.

- (b) Find the probability of a coincidence detection as function of the two-photon contamination