

II

Note: This problem covers the generation of coherent states based on the radiation from a classical dipole. Completing it is voluntary; you are encouraged to attempt it and should at least check out the 'official' solution set below. Whatever you choose to do, the problem will not count towards your homework score.

Consider a single mode of the electromagnetic field, coupled to a classical dipole $d(t)$ located at $z = 0$. As discussed in class, the Hamiltonian for this system can be written as

$$\hat{H}(t) = \hbar\omega(\hat{a}^\dagger\hat{a} + 1/2) + \hbar\lambda(t)(\hat{a} + \hat{a}^\dagger), \quad \lambda(t) = -\frac{d(t)\mathcal{E}_k}{\hbar}$$

In the following, work in the Schrödinger picture.

- (a) Calculate the commutators of \hat{a} and \hat{a}^\dagger with $\hat{H}(t)$.
- (b) Let $\alpha(t) = \langle \psi(t) | \hat{a} | \psi(t) \rangle$, where $|\psi(t)\rangle$ is the normalized state vector for the field. Show from the results of (a) that

$$\frac{d}{dt}\alpha(t) = -i\omega\alpha(t) - i\lambda(t)$$

Integrate this differential equation. At time t , what are the mean values of the quadrature operators $\hat{X}(t)$ and $\hat{Y}(t)$?

- (c) Let $|\varphi(t)\rangle = [\hat{a} - \alpha(t)]|\psi(t)\rangle$, where $\alpha(t)$ has the value calculated in (b). Using the results from (a) and (b), show that

$$i\hbar\frac{d}{dt}|\varphi(t)\rangle = [\hat{H}(t) + \hbar\omega]|\varphi(t)\rangle$$

How does the norm of $|\varphi(t)\rangle$ vary with time?

- (d) Assuming that $|\psi(0)\rangle$ is an eigenvector of \hat{a} with eigenvalue $\alpha(0)$, show that $|\psi(t)\rangle$ is also an eigenvector of \hat{a} , and calculate its eigenvalue.
- (e) Assume that at $t = 0$ the field is in the vacuum state. The dipole radiates between times 0 and T and is then removed. When $t > T$, what is the evolution of the mean values $\langle \psi(t) | \hat{X}(t) | \psi(t) \rangle$ and $\langle \psi(t) | \hat{Y}(t) | \psi(t) \rangle$?
- (f) Assume that between 0 and T the dipole oscillates so that $\lambda(t) = \lambda_0 \cos(\omega't)$. Derive an expression for $\alpha(T)$ as a function of $\Delta\omega = \omega' - \omega$. Sketch the variation of $|\alpha(T)|^2$ versus $\Delta\omega$. Discuss.

(Adapted from Cohen-Tannoudji's "Quantum Mechanics", Problem 6, p. 636)

A bit of helpful math for Problem III

The linear first-order differential equation

$$\frac{dy}{dx} + p(x)y = q(x)$$

has the solution

$$y(x) = e^{-P(x)} \left[\int_{x_0}^x e^{P(x')} q(x') dx' + y(x_0) e^{P(x_0)} \right]$$

where $P(x) = \int p(x) dx$ is the indefinite integral and $y(x_0)$ is the boundary value required to uniquely specify the solution y . The factor $e^{P(x_0)}$ ensures proper behavior for $q(x) = 0$. That this solution is valid is easy to check. For a derivation, see e. g. Arfken, Mathematical Methods for Physicists.

In problem III, substitute $x \rightarrow t$, $y(x) \rightarrow \alpha(t)$, $p(x) \rightarrow i\omega \Rightarrow e^{-P(x)} = e^{-i\omega t}$ and $q(x) \rightarrow i\lambda(t)$ to get

$$\alpha(t) = e^{-i\omega t} \left[\int_{t_0}^t e^{i\omega t'} i\lambda(t') dt' + \alpha(t_0) e^{i\omega t_0} \right]$$